# 5P7106 BMR

# BROWN MID-RIB SUDANGRASS X SUDANGRASS HYBRID

## **KEY POSITIONING / BENEFITS**

- Photoperiod Sensitive (Headless)
- **Extended Harvest Window**
- Best for Hay, Haylage & Grazing
- Tolerates High Cutting Frequencies Adapts Well to Intensive Grazing .
- Fine Stemmed & Leafy .

# **POSITIONING / MANAGEMENT**

#### SUMMER ANNUAL MANAGEMENT

Planting Date: Late April through mid August. Soil temperature should be 60°F to 65°F or warmer. Planting too early can cause slow early growth and reduced populatión.

Planting Depth: 3/4" to 1 1/2" deep depending on soil moisture.

Row Width: Drilled rows for best quality.

#### Planting Method: Drill preferred.

**Fertility:** Nitrogen: 75 to 100 lbs./acre PPI & 50 to 75 lbs./acre after first cutting. Phosphorus: 20 to 60 lbs./acre, more if soil test indicates less than 30 lbs./acre. Potassium: 40 to 80 lbs./acre, more if soil test indicates less than 300 lbs./acre.

#### PLANTING RATE (LBS./ACRE)

Principle	Planting	Average Annual Moisture (inches)		
Uses	Method	<20	20 - 26	26 - 32>
		Stress	Favorable	Irrigated <sup>2</sup>
	36" - 40" Rows	<b>4 - 8</b> <sup>1</sup>	8 - 12	12 - 15
Grazing & Cover Crop	20" - 30" Rows	<b>6 - 10</b> <sup>1</sup>	10 - 15	15 - 25
	Drilled	<b>10 - 15</b> <sup>1</sup>	15 - 20	20 - 30
Hay, Haylage & Greenchop	Drilled	15 - 20	20 - 25	25 - 50
Hay, Grazing & Cover Crop	Broadcast	<b>20 - 25</b> <sup>1</sup>	25 - 30	30 - 50

Recommended rates for cover crop
Includes high rainfall areas and conditions Planting rates for hybrid sudangrass and sorghum x sudangrass depends on a number of factors including location, growing conditions, management, etc. The above rates serve as a guide.

# AGRONOMIC TRAITS<sup>1</sup>

Forage Yield Potential	2
Early Growth Rate	2
Regrowth	1
Average Plant Height (feet)	7' - 9'
Standability	3
Drought Tolerance	
Leafiness	1
Avg. Seeds / Lb. (x1000)	25
Prussic Acid (HCN) Potential	

#### **DISEASE RESISTANCE**

MDMV Tolerance	Undetermined
Downy Mildew:	
Pathotype 1	Undetermined
Pathotype 3	Undetermined
Pathotype P6	Undetermined
Anthracnose	Undetermined

#### **INSECT RESISTANCE** Greenbug:

indug.	
Biotype C	Undetermined
Biotype E	Undetermined

### **PRINCIPAL USES<sup>1</sup>**

Hay	
Haylage	
Grazing / Pasture	2
Greenchop	4

# HARVEST MANAGEMENT

Do not cut or graze until forage reaches 18" in height. Cut at 24" to 36" for best quality and yield. Cut 6" to 8" above ground level for fastest regrowth. Condition with a crimper at a slight diagonal to speed drying and regrowth. Wilt to 50% to 60% moisture for storage in a bunker or concrete stave silo and 45% to 55% for a sealed storage system. Forage should be about 85% to 90% plant moisture at cutting. Chop at 1/4" to 1/8" cutting length to improve packing.

# **HARVEST HEIGHT**

Grazing	
Hay / Haylage	24" - 36"
Greenchop	

#### **YIELD**

2 to 4 cuttings depending on planting date, management and growing season. 3 to 6 tons/acre dry matter (20 to 40 tons green weight)

> Footnotes: 1 - Numerical Rating: 1 to 9 (1 = Excellent, 5 = Average, 9 = Poor) This brochure is for your information only. Please note that the ratings and this brock the S for your micromation only. Please note that the radings and descriptions of our products are based on research and field observations from multiple locations and years and may not be entirely consistent with the products that you will ultimately purchase from us. The sale of our products is subject to any agreement that we may have with you and to our terms and conditions as they contain important conditions of sale. terms and conditions as they contain important conditions of sale. including limitations of warranty and remedy. ©Copyright 2014. All rights reserved.

Sorghum Partners<sup>®</sup> word mark and logo are trademarks of S&W Seed Company or one of its affiliates



